Gamma rays from the littlest galaxies

Tracing Accretion and Dark Matter in the Low-Mass Universe

Milena Crnogorčević Postdoctoral Fellow Oskar Klein Centre/Stockholm University


ECAP Seminar October 23, 2025












Dwarf galaxies

→ Low-mass galaxies:

tens of $M_{sun} < M_{stellar} < 3 \times 10^9 M_{sun}$

- → At the forefront of current/future observational efforts:
- Near-field cosmology (Gaia, DES, Rubin/LSST)
- Early Universe & Reionization (JWST)
- Accretion & Feedback (Athena, AXIS, SKA, ngVLA)
- Fundamental physics & dark matter (Fermi, IACTs, IceCube)



Representative sample of the Local Group dwarf galaxies.


Dwarf galaxies

→ Low-mass galaxies:

tens of $M_{sun} < M_{stellar} < 3 \times 10^9 M_{sun}$


- → At the forefront of current/future observational efforts:
- Near-field cosmology (Gaia, DES, Rubin/LSST)
- Early Universe & Reionization (JWST)
- Accretion & Feedback (Athena, AXIS, SKA, ngVLA)
- Fundamental physics & dark matter (Fermi, IACTs, IceCube)

<u>Dwarf galaxies</u>: bridges between large scale structure, galaxy formation, & fundamental physics.



Representative sample of the Local Group dwarf galaxies.


Dwarf galaxies, big questions



How did the first black holes form?


- ✓ preserve signatures of seed formation
- ✓ test lower limits of BH-galaxy scaling relations
- ✓ probe accretion physics in low-mass regime

Dwarf galaxies, big questions



How did the first black holes form?


- ✓ preserve signatures of seed formation
- ✓ test lower limits of BH-galaxy scaling relations
- ✓ probe accretion physics in low-mass regime



What is the nature of dark matter?


- ✓ high dark matter densities
- ✓ min. baryonic contamination in some types
- \checkmark test λ CDM predictions at smallest scales

Dwarf galaxies, big questions



How did the first black holes form?

- ✓ preserve signatures of seed formation
- ✓ test lower limits of BH-galaxy scaling relations
- ✓ probe accretion physics in low-mass regime



What is the nature of dark matter?


- ✓ high dark matter densities
- ✓ min. baryonic contamination in some types
 - \checkmark test λ CDM predictions at smallest scales




How do galaxies evolve?

- ✓ stress-test feedback models in shallow potentials
- ✓ preserve imprints of reionization and early enrichment
- ✓ challenge numerical simulations at resolution limits


Dwarf or not?





Dwarf or not?






Globular star cluster 47 Tucanae The SMC: dwarf galaxy Akira Fuji/Hubble

Classification of dwarfs... is complicated.

Historically: morphology




[Gilmore+ 2007, Willman & Strader 2012, McConnachie 2012, Lelli et al. 2016, Simon 2019]


Globular star cluster 47 Tucanae The SMC: dwarf galaxy Akira Fuji/Hubble

Classification of dwarfs... is complicated.

Historically: morphology



[Gilmore+ 2007, Willman & Strader 2012, McConnachie 2012, Lelli et al. 2016, Simon 2019]




Classification of dwarfs... is complicated.

Historically: morphology

Now:

- → dynamical M/L ratio (indicating DM content)
- → half-light radius
- → velocity dispersion measurements
- → stellar populations (age, metallicity)
- → relaxation time
- → tidal dwarfs + faint halo systems especially difficult

[Gilmore+ 2007, Willman & Strader 2012, McConnachie 2012, Lelli et al. 2016, Simon 2019]




Dwarf elliptical galaxies (dEs)

Dwarf irregular galaxies (dlrrs)

Dwarf spheroidal galaxies (dSphs)

Ultrafaint dwarf galaxies (UFDs)

Ultracompact dwarf galaxies (UCDs)



e.g. Reines et al. 2013, Baldassare et al. 2015, 2023


Dwarf elliptical galaxies (dEs)

Dwarf irregular galaxies (dlrrs)


Dwarf spheroidal galaxies (dSphs)

Ultrafaint dwarf galaxies (UFDs)


Ultracompact dwarf galaxies (UCDs)



Obligatory slide on Fermi Large Area Telescope



Obligatory slide on Fermi Large Area Telescope



individual γ rays convert into e⁺e⁻ pairs \rightarrow tracks (localization) & deposited energy

*Energy range

**Field of View

***Single photon angular resolution

Timing accuracy

20 MeV to > 300 GeV

2.4 sr (~1/5 of the whole sky)

< 1 deg at 1 GeV

1 microsecond


*ideally suited for WIMP searches

**whole sky every ~3 hours

***point-source localization < 0.5 arcmin

...it also detects electrons.


Dwarf AGN




Based on Crnogorčević+ 2025, arXiv: **2509.18239**

Dwarf AGN

- \rightarrow host intermediate-mass black holes (IMBHs), $10^4 10^6 \, \mathrm{M}_{\odot}$
- → missing link in BH mass spectrum
- → preserve signatures of BH seed formation
- → enhanced dark matter density around IMBHs (potential dark matter targets)
- → but, multiple potential **γ**-ray production mechanisms...






Gammas from AGNs

Classical ways to produce γ -rays:

- → Accretion-related processes: Inverse Compton from corona/disk
- → Misaligned jets: Reduced Doppler boosting → softer spectra
- → Cosmic-ray interactions: Star formation/supernova activity
- → AGN-driven outflows: Shocks in interstellar medium

New ways to produce γ -rays:

→ Dark matter annihilation: Enhanced density around IMBHs



[Beckmann & Shrader 2013]

Multiple mechanisms likely contribute: requires multiwavelength approach to disentangle

→ Do gamma-ray emission mechanisms observed in massive galaxy AGN scale down to IMBHs?

- → Do gamma-ray emission mechanisms observed in massive galaxy AGN scale down to IMBHs?
- → How does emission efficiency depend on e.g., black hole mass as a proxy for accretion rate?

- → Do gamma-ray emission mechanisms observed in massive galaxy AGN scale down to IMBHs?
- → How does emission efficiency depend on e.g., black hole mass as a proxy for accretion rate?
- → Can gamma-ray observations distinguish between competing emission models?

- → Do gamma-ray emission mechanisms observed in massive galaxy AGN scale down to IMBHs?
- → How does emission efficiency depend on e.g., black hole mass as a proxy for accretion rate?
- → Can gamma-ray observations distinguish between competing emission models?
- → Are there **unique signatures** in IMBH systems not seen in SMBH-hosted AGN?

MaNGA AGN dwarf galaxies (MAD) - I. A new sample of AGN in dwarf galaxies with spatially resolved spectroscopy

M. Mezcua^{1,2★}, H. Domínguez Sánchez³


- ¹ Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Magrans, 08193 Barcelona, Spain
- ² Institut d'Estudis Espacials de Catalunya (IEEC), Carrer Gran Capità, 08034 Barcelona, Spain
- 3 Centro de Estudios de Física del Cosmos de Aragón (CEFCA), Plaza San Juan, 1, 44001, Teruel, Spain

Accepted 2024 January 24. Received 2024 January 8; in original form 2023 May 19

ABSTRACT

The finding of active galactic nuclei (AGN) in dwarf galaxies has important implications for galaxy evolution and supermassive black hole formation models. Yet, how AGN in dwarf galaxies form is still debated, in part due to scant demographics. We make use of the MaNGA survey, comprising $\sim 10,000$ galaxies at z < 0.15, to identify AGN dwarf galaxies using a spaxel by spaxel classification in three spatially-resolved emission line diagnostic diagrams (the [NII-, [SII]- and [OI]-BPT) and the WHAN diagram. This yields a sample of 664 AGN dwarf galaxies, the largest to date, and an AGN fraction of ~ 20% that is significantly larger than that of single-fiber-spectroscopy studies (i.e. $\sim 1\%$). This can be explained by the lower bolometric luminosity (< 10⁴² erg s⁻¹) and accretion rate (sub-Eddington) of the MaNGA AGN dwarf galaxies. We additionally identify 1,176 SF-AGN (classified as star-forming in the [NII]-BPT but as AGN in the [SII]- and [OI]-BPT), 122 Composite, and 173 LINER sources. The offset between the optical center of the galaxy and the median position of the AGN spaxels is more than 3 arcsec for ~62% of the AGN, suggesting that some could be off-nuclear. We also identify seven new broad-line AGN with log $M_{\rm BH} = 5.0 - 5.9 \ M_{\odot}$. Our results show how integral-field spectroscopy is a powerful tool for uncovering faint and low-accretion AGN and better constraining the demographics of AGN in dwarf galaxies.

Key words: Galaxies: dwarf, active, accretion



MaNGA AGN dwarf galaxies (M' DRAFT VERSION FEBRUARY 20, 2025
Typeset using latex two column
Typeset DRAFT VERSION FEBRUARY 20, 2025
Typeset using TATEX two column styl. dwarf galaxies with spatia¹¹ M. Mezcua^{1,2★}, H ⁻ 1 Institute of Space Scien ² Institut d'Estud; 3 Centro ' ACAUREPIKA FUCHA HVIDING Accepted 2024 n. HUNSCHEID, W. F. PRADA, R. MIQUEL, R. MIQUEL, ep nostic dia study black inte scaling relation a bette Which enable

Key wc

MOUSTAKAS, CIRCOSTA, BIANCHI, 14
MOUSTAKAS, CIRCOSTA, BIANCHI, 14
SY, AHLEN, 26,27 P. DOEL, 31

Draft version February 4, 2025 Typeset using IATEX twocolumn style in AASTeX631

the galaxy mass.

X-Ray Bright Active Galactic Nuclei in Local Dwarf Galaxies: Insights from eROSITA

Andrea Sacchi , ¹ Ákos Bogdán , ¹ Urmila Chadayammuri , ² and Angelo Ricarte ¹

¹Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 20138, USA ²Max Planck Institut für Astronomie, Königstuhl 17, 69121 Heidelberg, Germany

ABSTRACT

Although supermassive black holes (SMBHs) reside in the heart of virtually every massive galaxy, it remains debated whether dwarf galaxies commonly host SMBHs. Because low-mass galaxies may retain memory of the assembly history of their black holes (BHs), probing the BH occupation fraction of local dwarf galaxies might offer insights into the growth and seeding mechanisms of the first BHs. In this work, we exploit the Western half of the eROSITA all-sky survey (covering 20,000deg²) and compile a catalog of accreting SMBHs in local (D < 200 Mpc) dwarf galaxies. Cleaning our sample from X-ray background sources, X-ray binaries, and ultraluminous X-ray sources, we identify 74 AGN-dwarf galaxy pairs. Using this large and uniform sample, we derive a luminosity function of dwarf galaxy AGN, fitting it with a power law function and obtaining $dN/dL_X = (15.9 \pm 2.2) \times L_X^{-1.63 \pm 0.05}$. Measuring the offset between the dwarf galaxies centroids and the X-ray sources, we find that $\approx 50\%$ of the AGN are likely off-nuclear, in agreement with theoretical predictions. We compare the BH-to-stellar mass relation of our sample with the local and high-redshift relations, finding that our sources better adhere to the former, suggesting that local AGN across different mass scales underwent a similar growth history. Finally, we compare our sources with semi-analytical models: while our sample's shallowness prevents distinguishing between different seeding models, we find that the data favor models which keep SMBH in dwarf galaxies active at a moderate rate, motivating model improvement by comparison to AGN in the dwarf galaxy regime. DESI releases WIL Trevious low-reds

[astro-ph.GA]

Draft version February 4, 2025 Typeset using IATEX **twocolumn** style in AASTeX631

X-Ray Bright Active Galactic Nuclei in Local Dwarf Galaxies: Insights from eROSITA


Andrea Sacchi , ¹ Ákos Bogdán , ¹ Urmila Chadayammuri , ² and Angelo Ricarte , ¹

¹ Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 20138, USA
² Max Planck Institut für Astronomie, Königstuhl 17, 69121 Heidelberg, Germany


ABSTRACT

Although supermassive black holes (SMBHs) reside in the heart of virtually every massive galaxy, it remains debated whether dwarf galaxies commonly host SMBHs. Because low-mass galaxies may retain memory of the assembly history of their black holes (BHs), probing the BH occupation fraction of local dwarf galaxies might offer insights into the growth and seeding mechanisms of the first BHs. In this work, we exploit the Western half of the eROSITA all-sky survey (covering 20,000deg²) and compile a catalog of accreting SMBHs in local (D < 200 Mpc) dwarf galaxies. Cleaning our sample from X-ray background sources, X-ray binaries, and ultraluminous X-ray sources, we identify 74 AGN-dwarf galaxy pairs. Using this large and uniform sample, we derive a luminosity function of dwarf galaxy AGN, fitting it with a power law function and obtaining $dN/dL_X = (15.9 \pm 2.2) \times L_X^{-1.63 \pm 0.05}$. Measuring the offset between the dwarf galaxies centroids and the X-ray sources, we find that $\approx 50\%$ of the AGN are likely off-nuclear, in agreement with theoretical predictions. We compare the BH-to-stellar mass relation of our sample with the local and high-redshift relations, finding that our sources better adhere to the former, suggesting that local AGN across different mass scales underwent a similar growth history. Finally, we compare our sources with semi-analytical models: while our sample's shallowness prevents distinguishing between different seeding models, we find that the data favor models which keep SMBH in dwarf galaxies active at a moderate rate, motivating model improvement by comparison to AGN in the dwarf galaxy regime.

- → X-rays sensitive to accretion
- → minimizes the contamination or line emission biases
- → unbiased to obscuration of the AGN
- \rightarrow closest sources (D < 200 Mpc)
- → available control samples
- → almost... all-sky




Individual Source Analysis



Example, PGC 2077238

- 15 years of Fermi-LAT Pass 8
- Energy range: 500 MeV 500 GeV
- Event class: P8R3 SOURCE (evclass=128, evtype=3)
- Software: Fermipy v1.2.0 + Fermitools v2.2.0
- ROI: 10°×10° (model extended to 15°×15° for PSF spillover)
- Background sources from 4FGL-DR4
- Modeled as point source (unresolved at LAT resolution)
- Binned likelihood analysis per energy bin
- SED derived with fixed per-bin index $\Gamma = 2$

Individual Source Analysis: No Detection




Roughly, $\sqrt{TS} \sim$ detection significance (in σ). TS = 25 corresponds to \sim 5 σ detection in LAT.

- 15 years of Fermi-LAT Pass 8
- Energy range: 500 MeV 500 GeV
- Event class: P8R3 SOURCE (evclass=128, evtype=3)
- Software: Fermipy v1.2.0 + Fermitools v2.2.0
- ROI: 10°×10° (model extended to 15°×15° for PSF spillover)
- Background sources from 4FGL-DR4
- Modeled as point source (unresolved at LAT resolution)
- Binned likelihood analysis per energy bin
- SED derived with fixed per-bin index $\Gamma = 2$

What does this no detection mean?

X-ray / γ -ray Scaling

• Empirical **BL Lac relation**




 $\log F_{\gamma} = 0.42 \log F_X - 8.17$ (& scatter based on different states)


What does this no detection mean?

X-ray / γ -ray Scaling

• Empirical **BL Lac relation**




Most predicted fluxes fall just below LAT's sensitivity after 15 years



 $\log F_{V} = 0.42 \log F_{X} - 8.17$ (& scatter based on different states)

What does this no detection mean?

Most predicted fluxes fall just below LAT's sensitivity after 15 years



- No outliers detected → BL Lac-like SSC emission likely suppressed
- Dwarf AGN are ≥ 7x less efficient in producing gamma rays than BL Lacs (at fixed X-ray flux)
- Possible causes:
 - Weak or misaligned jets
 - Different accretion regimes
 - $\gamma\gamma$ absorption in denser environments

Joint Likelihood Analysis: Population-level Gamma-ray Signal


- → No significant individual detections
- → Use **joint-likelihood analysis**, summing log-likelihoods from all 67 sources (we excluded 7 sources due to their latitude/proximity to associated LAT sources)


$$\log L_{total} = \sum_{i} \log L_{i} \left(c_{i'} \Gamma \mid D_{i} \right)$$


- \rightarrow Common spectral shape (power law index Γ) across all sources
- ightharpoonup Source normalizations tied by physical weights w_i
- → Blank skies used for calibration


Advantages


- Retains background, exposure, and brightness differences
- Statistically robust compared to simple stacking











Interpretation:

- → Suggestive-but not conclusive-evidence that **γ**-ray emission may scale with IMBH mass
- \rightarrow Spectrum significantly softer than typical blazars ($\Gamma \approx 2-2.5$)
- → Possible explanations:
 - Coronal or accretion-disk Comptonization
 - Weak or misaligned jets
 - **yy** absorption or host star formation contamination
- → Verified against 400 blank-field ensembles and excluded-source control sample
- → Similar soft peak appears when extended down to 100 MeV (but with inflated significance → likely systematics)



dSphs


- → most dark-matter dominated systems known in the Universe
- → 61 dwarf satellites identified so far in the Milky Way, more on the horizon
- → nearby (furthest 460 kpc), some within single-digit kpc
- → not much astrophysical background, inactive




dSphs

DM γ -ray flux

 $\frac{d\Phi}{dE}$




astrophysics J-factor




Dark matter content determined from stellar velocity dispersion

- **Classical dwarfs**: spectra for several thousand stars
- **Ultra-faint dwarfs**: spectra for fewer than 100 stars




particle physics

$$\frac{\langle \sigma v \rangle}{2M_{DM}^2} \sum B_i \frac{dN_{\gamma}}{dE}$$




J-factors [Fermi-LAT Collaboration 2024]


Combined dSph Analyses



[Fermi-LAT Collaboration '17]




[Fermi-LAT Collaboration '21]



[McDaniel+ '24]

Shaded regions: blank-field analysis

Think: $\sqrt{TS} \sim \sigma$




Combined dSph Analyses

(2024)

ightharpoonup 7 dSphs with local significance > 2 σ


... not the first time: some marginal significance reports in the past.

Ret II (DiMauro+ '21, Albert+ '17, Geringer-Sameth+ '15, Hooper & Linden '15) Tucana II, Willman 1, Horologium II, Bootes I (DiMauro+ '21)



Think: $\sqrt{TS} \sim \sigma$

Limits on the parameter space




 \rightarrow Trials factor reduces significance to 0.5 σ .

Observations:

- → generally consistent with previous limits; in **tension** with the GCE results
- → cannot rule out DM due to the uncertainties in galactic center analysis (DM profile and interstellar diffuse model) and uncertainties in J-factor of dwarfs


J-value calculations are... hard


[McDaniel+ '24]



Many underlying assumptions: dark matter distribution models, parametric/non-parametric approaches, observational limitations – can result in at least a factor of few difference to the real value. [e.g., Bonnivard+ '15, Geringer-Sameth+ '15, Hayashi+ '16, Ando+ '20 etc.]

Future of dSph DM searches






[Fermi-LAT Collaboration '24]

- Perform multi-instrument and multi-target analysis to obtain the most sensitive and robust results
- Joining likelihoods across instruments is challanging
- Focus: dSphs
- Limits driven by LAT sensitivity
- Legacy analysis of the current-generation gamma-ray instruments















PREPARED FOR SUBMISSION TO JCAP


Combined dark matter search towards dwarf spheroidal galaxies with Fermi-LAT, HAWC, WILLIAM MAGIC, and VERITAS H.E.S.S., MAGIC, and VERITAS

The Fermi-LAT, HAWC, H.E.S.S., MAGIC, and VERITAS Collaborations























Dominated by Fermi LAT

1










HAWC, HESS, MAGIC, VERITAS take over

2

How many dwarf galaxies do we *really* need? Maybe just one, but a good one?

Ursa Major III




[MC & Linden '23]

Ursa Major III

[Discovery: Smith+ 2023]

[J-factor: Errani+ 2023]

- → Unstable unless large DM content
- \rightarrow Nearby (~10 kpc)
- → Strong constraints on DM annihilation
- → Confirming the dark matter density requires deeper optical surveys



No Observational Evidence for Dark Matter Nor a Large Metallicity Spread in the Extreme Milky Way Satellite Ursa Major III / UNIONS 1

WILLIAM CERNY, DAISY BISSONETTE, ALEXANDER P. JI, MARLA GEHA, ANIRUDH CHITI, SIMON E.T. SMITH, JOSHUA D. SIMON, ANDREW B. PACE, EVAN N. KIRBY, KIM A. VENN, TING S. LI, AND ALICE M. LUNA,

¹Department of Astronomy, Yale University, New Haven, CT 06520, USA

²Department of Astronomy and Astrophysics, University of Chicago, Chicago, IL 60637, USA

³Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA

⁴NSF-Simons AI Institute for the Sky (SkAI), 172 E. Chestnut St., Chicago, IL 60611, USA

⁵Department of Physics and Astronomy, University of Victoria, Victoria, BC, V8P 1A1, Canada

⁶Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA

⁷Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904 USA

⁸Department of Physics and Astronomy, University of Notre Dame, 225 Nieuwland Science Hall, Notre Dame, IN 46556, USAA

⁹Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto ON, M5S 3H4, Canada

ABSTRACT

The extremely-low-luminosity, compact Milky Way satellite Ursa Major III / UNIONS 1 (UMaIII/U1; $L_V = 11 \ L_\odot$; $a_{1/2} = 3$ pc) was found to have a substantial velocity dispersion at the time of its discovery ($\sigma_v = 3.7^{+1.4}_{-1.0} \ \text{km s}^{-1}$), suggesting that it might be an exceptional, highly dark-matter-dominated dwarf galaxy with very few stars. However, significant questions remained about the system's dark matter content and nature as a dwarf galaxy due to the small member sample (N = 11), possible spectroscopic binaries, and the lack of any metallicity information. Here, we present new spectroscopic observations covering N = 16 members that both dynamically and chemically test UMaIII/U1's true nature. From higher-precision Keck/DEIMOS spectra, we find a 95% confidence level velocity dispersion limit of $\sigma_v < 2.3 \ \text{km s}^{-1}$, with a ~ 120 :1 likelihood ratio favoring the expected stellar-only dispersion of $\sigma_* \approx 0.1 \ \text{km s}^{-1}$ over the original 3.7 km s⁻¹ dispersion. There is now no observational evidence for dark matter in the system. From Keck/LRIS spectra targeting the Calcium



Segue into conclusions

Dwarf AGN

- \rightarrow No individual detections in γ rays emission suppressed or shifted to softer spectra.
- \rightarrow Joint-likelihood hints at mass-dependent γ -ray scaling with IMBHs warrants deeper (preferably MeV) study.

dSphs


- → Combined analyses remain consistent with limits; hints persist but are statistically marginal.
- → Next-generation surveys (Rubin/LSST, DECam, etc.) will expand the dSph census and improve sensitivity.

Looking forward


- \rightarrow Joint, multi-instrument efforts (LAT, HAWC, H.E.S.S., MAGIC, VERITAS) set the stage for a legacy analysis of the γ -ray sky.
- → The smallest galaxies may be our ticket to fame.








MeV GeV TeV PeV






IceCube (50 GeV - several PeV)





ANTARES (50 GeV - 100 TeV)



KM3NeT/ARCA (100 GeV - several PeV)



KM3NeT/ORCA (1 - 100 GeV)

 ν s (future)


IceCube-Gen2 In the works: 8 km³ + radio








IceCube (50 GeV - several PeV)



IceCube Upgrade

ANTARES (50 GeV - 100 TeV)



KM3NeT/ARCA (100 GeV - several PeV)



KM3NeT/ORCA (1 – 100 GeV)





GeV


TeV


PeV














- MIDEX and SMEX: Swift and NICER
- Technology: a robust technology development program (SiPMs, new scintillators, upgraded silicon detectors, etc)
 Balloons (+ CubeSats!): long duration balloons enabled
- COSI, LEAP, etc.
- Data Analysis & Theory: mainly supported through GI programs
- TeV Astronomy: VERITAS, HESS, HAWC, and MAGIC.



[insert your space-based gamma-ray wish list]



[insert your space-based gamma-ray wish list]




Submitted to the NASA Astrophysics Advisory Committee by The Future Innovations in Gamma Ray Science Analysis Group


Future Innovations in Gamma Rays SAG: A Report on Gamma-ray Science Objectives Beyond 2025

Chris Fryer¹, C. Michelle Hui², Paolo Coppi³, Milena Crnogorcevic⁴, Tiffany R. Lewis⁵, Marcos Santander⁶, and Zorawar Wadiasingh⁷

¹Los Alamos National Laboratory
 ²NASA Marshall Space Flight Center
 ³Yale University
 ⁴Stockholm University
 ⁵Michigan Technological University
 ⁶University of Alabama, Huntsville
 ⁷University of Maryland, College Park




- Overview purpose & limitations in scope
- Context state of the field & progress since the 2020 Decadal
- Key Science Opportunities that Require Gamma-ray Observations
 - Nuclear Decay requires high angular resolution & high sky coverage
 - GRBs need high sky coverage & fast alerts
 - Blazars need high effective area & polarization
 - Pulsar Timing Arrays need high precision timing and high angular resolution
 - Dark Matter
- Theory & computation are currently limited by lack of precision in observables
- Technology Advancement a lot has happened that has not flown
- Mission Capabilities & Infrastructure the success of missions requires that the surrounding infrastructure function appropriately.




JHEAP Special Issue


- Provides a place to document supporting work and credit authors of individual plots, studies and strategic arguments.
- Currently accepting manuscripts until 7 Nov
 2025 (likely extended)
- If you are intending to submit an article and have not been in touch with us, please reach out now.
- If you have an idea of something you would like to help us demonstrate, then please let us know.




1968, Orbiting Solar Observatory, OSO-3 (~50 MeV)



2000, COMPTEL (onboard CGRO), 1–30 MeV



2000, EGRET (onboard CGRO), above 100 MeV



2020, LAT (onboard *Fermi*), above 500 MeV

Physics of the Cosmos

Exploring fundamental questions regarding the physical forces of the universe

Due to the lapse in federal government funding, NASA is not updating this website. We sincerely regret this inconvenience.

FIG SAG Home

SAG Leadership

SAG Events

Future Innovations in Gamma Rays (FIG SAG)

Website https://pcos.gsfc.nasa.gov/sags/figsag.php

Slack Workspace https://docs.google.com/forms/d/e/1FAlpQLSfsgnb1OUQ3jlSGil

M 3abQsKoHvzlgWBZP3meMXJxUwRHI5w/viewform

Chairs e-mail addresses Michelle Hui (c.m.hui@nasa.gov)

Chris Fryer (<u>fryer@lanl.gov</u>)

Co-chairs e-mail addresses Paolo Coppi (paolo.coppi@yale.edu)

Milena Crnogorčević (milena.crnogorcevic@fysik.su.se)

Tiffany Lewis (tiffanylewisphd@gmail.com)

Zorawar Wadiasingh (<u>zorawar.wadiasingh@nasa.gov</u>)

News

22 September 2025

Final Reminder: PhysCOS Second Early Career Workshop on 23-25 September 2025

» Details